



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11) Publication number:

**0 352 960 B1**

(12)

## EUROPEAN PATENT SPECIFICATION

- (45) Date of publication of patent specification: **26.10.94** (51) Int. Cl.<sup>5</sup>: **C07D 473/30, C07D 239/48C3, A61K 31/52**
- (21) Application number: **89307341.1**
- (22) Date of filing: **19.07.89**

(54) **Purine compounds, process for their preparation and pharmaceutical compositions.**

- (30) Priority: **25.07.88 GB 8817651**
- (43) Date of publication of application:  
**31.01.90 Bulletin 90/05**
- (45) Publication of the grant of the patent:  
**26.10.94 Bulletin 94/43**
- (84) Designated Contracting States:  
**AT BE CH DE ES FR GB GR IT LI LU NL SE**
- (56) References cited:  
**EP-A- 0 293 063**  
**GB-A- 1 338 235**

- (73) Proprietor: **SMITH KLINE & FRENCH LABORATORIES LIMITED**  
**Mundells**  
**Welwyn Garden City Hertfordshire, AL7 1EY (GB)**
- (72) Inventor: **Coates, William John**  
**SmithKline Beecham Pharmaceuticals, The Frythe**  
**Welwyn, Hertfordshire AL6 9AR (GB)**
- (74) Representative: **Waters, David Martin, Dr. et al**  
**SmithKline Beecham plc**  
**Corporate Intellectual Property**  
**SB House**  
**Great West Road**  
**Brentford, Middlesex TW8 9BD (GB)**

**EP 0 352 960 B1**


Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

## Description

GB-A-1338235 discloses 8-azapurin-6-one derivatives useful in the treatment of respiratory disorders manifested by the interaction of tissue-fixed antibodies with specific antigens, such as allergic bronchial asthma.

The present invention relates to purinone derivatives and in particular to such compounds having a substituted phenyl group at the 2-position of the purinone ring. This invention further relates to processes for their preparation, intermediates in their preparation, their use as therapeutic agents and to pharmaceutical compositions containing them. The compounds of this invention are inhibitors of a calmodulin insensitive cyclic GMP phosphodiesterase and are of use in combatting such conditions where such inhibition is thought to be beneficial. They are bronchodilators and are therefore of use in combatting chronic reversible obstructive lung diseases such as asthma and bronchitis. Some of the compounds of the present invention have anti-allergic activity and are therefore useful in combatting allergic diseases such as allergic asthma, allergic rhinitis, urticaria and irritable bowel syndrome. Furthermore the compounds of this invention are vasodilators and are therefore of value in combatting angina, hypertension and congestive heart failure.

Accordingly the present invention provides compounds of the formula (1) :



and pharmaceutically acceptable salts thereof, wherein

- R<sup>1</sup> is C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>3-5</sub> cycloalkyl, C<sub>1-4</sub> alkyl, phenyl, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkyl substituted by 1 to 6 fluoro groups;
- R<sup>2</sup> is hydrogen, hydroxy, C<sub>1-4</sub> alkyl, phenyl, mercapto, C<sub>1-4</sub> alkylthio, CF<sub>3</sub> or amino;
- R<sup>3</sup> is hydrogen, nitro, amino, C<sub>1-4</sub> alkanoylamino, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> alkyl, halo, SO<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>, CONR<sup>4</sup>R<sup>5</sup>, cyano or C<sub>1-4</sub> alkylS(O)<sub>n</sub>;
- R<sup>4</sup> and R<sup>5</sup> are independently hydrogen or C<sub>1-4</sub> alkyl; and
- n is 0, 1 or 2;

provided that R<sup>3</sup> is not hydrogen when R<sup>1</sup> is C<sub>1-6</sub> alkyl or C<sub>2-6</sub> alkenyl and R<sup>2</sup> is hydrogen or hydroxy.

Suitably R<sup>1</sup> is C<sub>2-5</sub> alkyl for example ethyl, n-propyl, isopropyl, butyl, isobutyl or pentyl.

Suitably R<sup>1</sup> is C<sub>3-5</sub> alkenyl for example allyl, butenyl or pentenyl.

Suitably R<sup>1</sup> is cyclopropylmethyl.

Examples of C<sub>1-4</sub> alkyl substituted by 1 to 6 fluoro groups include -CF<sub>3</sub>, -CH<sub>2</sub>CF<sub>3</sub> or -CF<sub>2</sub>CHFCF<sub>3</sub>.

Preferably R<sup>1</sup> is n-propyl.

Suitably R<sup>2</sup> is hydrogen or hydroxy.

Suitably R<sup>2</sup> is phenyl or C<sub>1-4</sub> alkyl for example methyl, ethyl, propyl, or butyl.

Suitably R<sup>2</sup> is mercapto or C<sub>1-4</sub> alkylthio for example methylthio or ethylthio.

Suitably R<sup>3</sup> is hydrogen.

Suitably R<sup>3</sup> is nitro, cyano, CONR<sup>4</sup>R<sup>5</sup> or SO<sub>2</sub>NR<sup>4</sup>R<sup>5</sup> wherein NR<sup>4</sup>R<sup>5</sup> is amino, methylamino or dimethylamino.

Suitably R<sup>3</sup> is C<sub>1-4</sub> alkanoylamino, C<sub>1-4</sub> alkoxy or C<sub>1-4</sub> alkyl for example acetamido, methoxy, ethoxy, methyl or ethyl.

Suitably R<sup>3</sup> is halo for example fluoro, chloro, bromo or iodo.

Particular compounds of this invention are :

2-(2-[2,2,2-trifluoroethoxy]phenyl)purin-6-one,

2-(2-cyclopropylmethoxyphenyl)purin-6-one,

2-(2-cyclopropylmethoxyphenyl)purin-6,8-dione,

2-(2-benzyloxyphenyl)purin-6,8-dione,


2-(2-propoxyphenyl)-8-trifluoromethylpurin-6-one,

2-(2-propoxyphenyl)-8-phenylpurin-6-one,

2-(2-propoxyphenyl)-8-methylpurin-6-one,

2-(2-propoxyphenyl)-8-mercaptapurin-6-one,  
 2-(2-propoxyphenyl)-8-methylthiopurin-6-one,  
 2-(2-propoxyphenyl)-8-aminopurin-6-one,  
 2-(2-propoxy-5-nitrophenyl)purin-6-one,  
 5 2-(2-propoxy-5-aminophenyl)purin-6-one,  
 2-(2-propoxy-5-acetamidophenyl)purin-6-one,  
 2-(2-propoxy-4-methoxyphenyl)purin-6-one,  
 2-(2-propoxy-5-methoxyphenyl)purin-6-one,  
 2-(2-propoxy-5-chlorophenyl)purin-6-one,  
 10 2-(2-propoxy-4-methylphenyl)purin-6-one,  
 2-(2-propoxy-5-fluorophenyl)purin-6-one,  
 2-(2-propoxy-5-dimethylsulphamoylphenyl)purin-6-one,  
 2-(2-propoxy-5-methylsulphamoylphenyl)purin-6-one,  
 2-(2-propoxy-5-sulphamoylphenyl)purin-6-one,  
 15 2-(2-propoxy-4-methylthiophenyl)purin-6-one,  
 2-(2-propoxy-5-cyanophenyl)purin-6-one, or  
 2-(2-propoxy-5-carbamoylphenyl)purin-6-one,  
 or pharmaceutically acceptable salts thereof.

This invention covers all tautomeric and optical isomeric forms of compounds of formula (1). For  
 20 example the compound of the formula (1) wherein R<sup>2</sup> is hydroxy or mercapto can exist in a tautomeric form



wherein X is O or S.

Compounds of the formula (1) wherein R<sup>2</sup> is hydrogen or amino, or R<sup>3</sup> is amino may form pharmaceuti-  
 35 cally acceptable salts with acids such as hydrochloric, hydrobromic, sulphuric, phosphoric, acetic, citric,  
 maleic, lactic, ascorbic, fumaric, oxalic, methanesulphonic and ethanesulphonic acids.

Compounds of the formula (1) may form pharmaceutically acceptable salts with metal ions, such as  
 alkali metals for example sodium and potassium, or with an ammonium ion.

In order to use a compound of the formula (1) or a pharmaceutically acceptable salt thereof for the  
 40 treatment of humans and other mammals it is normally formulated in accordance with standard pharmaceu-  
 tical practice as a pharmaceutical composition.

Compounds of formula (1) and their pharmaceutically acceptable salts may be administered in standard  
 manner for the treatment of the indicated diseases, for example orally, sublingually, parenterally, transder-  
 mally, rectally, via inhalation or via buccal administration.

45 Compounds of formula (1) and their pharmaceutically acceptable salts which are active when given  
 orally or via buccal administration can be formulated appropriately in dosage forms such as liquids, syrups,  
 tablets, capsules and lozenges. An oral liquid formulation will generally consist of a suspension or solution  
 of the compound or salt in a liquid carrier for example, ethanol, glycerine or water with a flavouring or  
 colouring agent. Where the composition is in the form of a tablet, any pharmaceutical carrier routinely used  
 50 for preparing solid formulations may be used. Examples of such carriers include starch, celluloses, lactose,  
 sucrose and magnesium stearate. Where the composition is in the form of a capsule, any routine  
 encapsulation process may be suitable, for example using the aforementioned carriers in a hard gelatin  
 capsule shell. Where the composition is in the form of a soft gelatin shell capsule, any pharmaceutical  
 carrier routinely used for preparing dispersions or suspensions may be considered, for example aqueous  
 55 gums, celluloses, silicates or oils and are incorporated in a soft gelatin capsule shell.

Typical parenteral compositions consist of a solution or suspension of the compound or salt in a sterile  
 aqueous or non-aqueous carrier optionally containing a parenterally acceptable oil or solubilising agent, for  
 example polyethylene glycol, polyvinylpyrrolidone, 2-pyrrolidone, cyclodextrin, lecithin, arachis oil or ses-

ame oil.

A typical suppository formulation comprises a compound of formula (1) or a pharmaceutically acceptable salt thereof which is active when administered in this way, with a binding and/or lubricating agent, for example polymeric glycols, gelatins, cocoa-butter or other low melting vegetable waxes or fats or their synthetic analogues.

Typical transdermal formulations comprise a conventional aqueous or non-aqueous vehicle, for example a cream, ointment, lotion or paste or are in the form of a medicated plaster, patch or membrane.

Typical compositions for inhalation are in the form of a solution, suspension or emulsion that may be administered in the form of an aerosol using a conventional propellant such as dichlorodifluoromethane or trichlorofluoromethane, or are in the form of a powder for insufflation.

Preferably the composition is in unit dosage form, for example a tablet, capsule or metered aerosol dose, so that the patient may administer to himself a single dose.


Each dosage unit for oral administration contains suitably from 0.001 mg/Kg to 30 mg/Kg, and preferably from 0.005 mg/Kg to 15 mg/Kg, and each dosage unit for parenteral administration contains suitably from 0.001 mg/Kg to 10 mg/Kg, of a compound of formula (1) or a pharmaceutically acceptable salt thereof calculated as the free base.

The daily dosage regimen for oral administration is suitably about 0.001 mg/Kg to 120 mg/Kg, of a compound of formula (1) or a pharmaceutically acceptable salt thereof calculated as the free base. The daily dosage regimen for parenteral administration is suitably about 0.001 mg/Kg to 40 mg/Kg, for example about 0.005 mg/Kg to 10 mg/Kg, of a compound of the formula (1) or a pharmaceutically acceptable salt thereof calculated as the free base. The active ingredient may be administered as required, for example from 1 to 8 times a day or by infusion. The compositions of the invention are bronchodilators and are useful in chronic reversible obstructive lung disease for example asthma and bronchitis. Some of the compositions of the present invention have anti-allergic activity and are therefore useful in combating allergic diseases such as allergic asthma, allergic rhinitis, urticaria and irritable bowel syndrome. The compositions of the present invention have vasodilator activity and are of use in the treatment of angina, hypertension and congestive heart failure. Such conditions can be treated by administration orally, sublingually, topically, rectally, parenterally or by inhalation. For administration by inhalation dosages are controlled by a valve, are administered as required and for an adult are conveniently in the range 0.1-5.0 mg of a compound of the formula (1) or a pharmaceutically acceptable salt thereof.

The compounds of this invention may be co-administered with other pharmaceutically active compounds, for example in combination, concurrently or sequentially. Conveniently the compounds of this invention and the other active compound or compounds are formulated in a single pharmaceutical composition. Examples of compounds which may be included in pharmaceutical compositions with the compounds of the formula (1) are bronchodilators such as sympathomimetic amines for example isoprenaline, isoetharine, salbutamol, phenylephrine and ephedrine or xanthine derivatives for example theophylline and aminophylline, anti-allergic agents for example disodium cromoglycate, histamine H<sub>1</sub>-antagonists, vasodilators for example hydralazine, angiotensin converting enzyme inhibitors for example captopril, anti-anginal agents for example isosorbide nitrate, glyceryl trinitrate and pentaerythritol tetranitrate, anti-arrhythmic agents for example quinidine, procainamide and lignocaine, calcium antagonists for example verapamil and nifedipine, diuretics such as thiazides and related compounds for example bendroflumazide, chlorothiazide, chlorothalidone, hydrochlorothiazide, and other diuretics for example frusemide and triamterene, and sedatives for example nitrazepam, flurazepam and diazepam.

In another aspect the present invention provides a process for the preparation of a compound of the formula (1) or a pharmaceutically acceptable salt thereof, which process comprises:

- a) for compounds wherein R<sup>2</sup> is hydrogen, C<sub>1</sub>-4-alkyl, phenyl or CF<sub>3</sub>, reacting a compound of the formula (2):



(2)

wherein  $R^1$  is as hereinbefore defined and  $R^6$  is a group  $R^3$  as hereinbefore defined or a precursor thereof,

with a compound of the formula  $R^7COL$  or a chemical equivalent thereof wherein  $R^7$  is hydrogen,  $C_1-4$  alkyl, phenyl or  $CF_3$  and  $L$  is a leaving group;

5 b) for compounds wherein  $R^2$  is hydroxy,

reacting a compound of the formula (2) as hereinbefore defined with a carbonylating agent;

c) for compounds wherein  $R^2$  is mercapto,

reacting a compound of the formula (2) as hereinbefore defined with thiourea;

d) for compounds wherein  $R^2$  is  $C_1-4$  alkylthio,

10 reacting a compound of the formula (2) as hereinbefore defined with thiourea and thereafter with a  $C_1-4$  alkyl halide;

e) for compounds wherein  $R^2$  is amino,

reacting a compound of the formula (2) as hereinbefore defined with cyanogen bromide;

and thereafter where necessary :

15 • converting a group  $R^6$  to a group  $R^3$ ;

• optionally forming a pharmaceutically acceptable salt.

The reaction between a compound of the formula (2) and a compound of the formula  $R^7COL$  is conveniently performed in the absence of a solvent or in the presence of a suitable solvent such as a  $C_1-4$  alcohol, pyridine or  $N$ -methylpyrrolidone, at ambient or elevated temperature, for example  $50-250^\circ C$ ,  
20 preferably  $100-200^\circ C$ . Suitably  $L$  is selected from hydroxy,  $C_1-4$  alkoxy, halo such as chloro or bromo, amino,  $C_1-4$  alkylamino or  $OCOR^7$  thus forming an acid anhydride. By a chemical equivalent of  $R^7COL$  is meant a reagent that will react in similar manner with a compound of the formula (2) to form a purine ring. Examples include amidines of the formula  $R^7C(NH)L^1$  wherein  $L^1$  is amino or  $C_1-4$  alkylamino, and alkyl orthoformates of the formula  $R^7C(L^2)_3$  wherein  $L^2$  is  $C_1-4$  alkoxy.

25 The reaction between a compound of the formula (2) and a carbonylating agent is conveniently performed in the absence of a solvent or in a suitable solvent such as a halohydrocarbon, pyridine or toluene, at ambient or elevated temperature, for example  $50-250^\circ C$ . Suitable carbonylating agents include urea,  $di(C_1-4)alkyl$ -carbonate,  $C_1-4$  alkyl chloroformate, phosgene, trichloromethyl chloroformate or carbonyldiimidazole.

30 The reaction between a compound of the formula (2) and thiourea is conveniently performed in the absence of a solvent or in a suitable solvent such as a halohydrocarbon, pyridine or toluene, at elevated temperature for example  $50-250^\circ C$ , optionally in the presence of a base such as potassium acetate. If desired the product from this reaction can suitably be reacted with a  $C_1-4$  alkyl halide, for example methyl iodide, in the presence of a base such as aqueous sodium hydroxide at ambient or elevated temperature  
35 e.g.  $40-100^\circ C$ .

A compound of the formula (2) is suitably reacted with cyanogen bromide in a solvent such as a  $C_1-4$  alkanol, pyridine or acetonitrile optionally in the presence of water at ambient or elevated temperature, e.g.  $40-100^\circ C$ .

40 Examples of  $R^6$  being a precursor to a group  $R^3$  is when  $R^6$  is a  $C_1-4$  alkylthio or  $C_1-4$  alkoxy carbonyl group.

Suitably a  $C_1-4$  alkylthio group can be converted to a  $C_1-4$  alkylsulphinyl group by treatment with one mole equivalent of an oxidising agent such as hydrogen peroxide or potassium periodate. A further mole equivalent of an oxidising agent such as hydrogen peroxide or potassium permanganate can be used to convert a  $C_1-4$  alkylsulphinyl group to a  $C_1-4$  alkylsulphonyl group.

45 A  $C_1-4$  alkoxy carbonyl group can be converted to a  $CONR^4R^5$  group by reaction with an amine  $HNR^4R^5$  wherein  $R^4$  and  $R^5$  are as hereinbefore defined.

A compound of the formula (1) wherein  $R^3$  is hydrogen can be converted to the corresponding compound wherein  $R^3$  is 5-nitro by reaction with a suitable nitrating agent, such as fuming nitric acid together with sulphuric acid.

50 A compound of the formula (1) wherein  $R^3$  is nitro can be converted to the corresponding compound wherein  $R^3$  is amino by reaction with a reducing agent, for example via catalytic hydrogenation with palladium on carbon. If desired the amino group can be converted to a  $C_1-4$  alkanoylamino group by treatment with a  $C_1-4$  alkanoylating agent, for example acetic anhydride.

Alternatively a compound of the formula (1) wherein  $R^3$  is amino can be treated with sodium nitrite and  
55 an inorganic acid such as sulphuric acid to form a diazonium salt which can be converted to compounds of the formula (1) wherein  $R^3$  is cyano or halo by reaction with cuprous cyanide or cuprous halide.

A compound of the formula (1) wherein  $R^3$  is cyano can be hydrolysed to the corresponding compound wherein  $R^3$  is carboxamido by treatment with concentrated sulphuric acid or by treatment with hydrogen

peroxide and potassium hydroxide.

A compound of the formula (1) wherein  $R^3$  is hydrogen can be converted to the corresponding compound wherein  $R^3$  is  $5-SO_2NR^4R^5$  by reaction with chlorosulphonic acid and thereafter with an amine  $HNR^4R^5$  wherein  $R^4$  and  $R^5$  are as hereinbefore defined.

As will be readily understood by the man skilled in the art, reactive groups in other parts of the molecule (e.g. when  $R^2$  is amino) can be protected before the group  $R^3$  is converted as hereinbefore described.

Compounds of the formula (2) are known or preparable in conventional manner from US Patents 3819631 and 4039544.

Pharmaceutically acceptable acid addition salts of the compounds of the formula (1) wherein  $R^2$  is hydrogen or amino, or  $R^3$  is amino may be prepared from the corresponding base of the compounds of the formula (1) in conventional manner. For example the base may be reacted with an acid in a  $C_1-4$  alkanol, or an ion-exchange resin may be used. The salts of the compounds of the formula (1) may be interconverted using ion-exchange resins. Non-pharmaceutically acceptable salts are therefore of use as they can be converted to pharmaceutically acceptable salts.

Pharmaceutically acceptable base addition salts of the compounds of the formula (1) may be prepared by standard methods, for example by reacting a solution of the compound of the formula (1) with a solution of the base.

The following biological test method, data, descriptions and Examples serve to illustrate this invention.

#### Bronchodilatation - In vivo

Male guinea-pigs of the Dunkin Hartley strain (500 - 600g) were anaesthetised with Sagatal (pentobarbital sodium) (60 mg/kg). Airway resistance was measured using a modification of the classical Konzett-Rossler technique (J. Pharm. Methods, 13, 309-315, 1985). U46619 (9,11-methanoepoxy-PGH<sub>2</sub>) was infused i.v. at a rate of 2.5 nmol/min, this produced a steady state of bronchoconstriction (approximately 120% increase from basal airway resistance). The compound under test was administered by i.v. bolus injection, and the subsequent peak inhibition of bronchoconstriction recorded.

The dose of compound required to reduce the U46619-induced bronchoconstriction by 50% is given as the  $BD_{50}$ . The compounds of Examples 2, 3, 6, 11, 14 and 15 had  $BD_{50}$  values in the range 0.58-7.24  $\mu$ mol/kg. These results demonstrate in vivo anti-bronchoconstrictor activity.

#### Anti-allergic activity

Male Duncan Hartley guinea-pigs (250-300 g) were sensitised to ovalbumen by i.p. injection of 2 ml of 50mg.ml<sup>-1</sup> i.p. and 0.2 ml s.c. Three weeks later they were anaesthetised with 60mg.kg<sup>-1</sup> sodium pentobarbitone. The trachea was cannulated and the animal respired at a rate of 40 breaths per minute and at an initial tracheal inflation pressure of 16 mmHg. Tracheal inflation pressure was measured by a transducer connected to a side arm of the respiration circuit. The carotid artery was cannulated for the measurement of blood pressure and the signal was used to trigger an instantaneous rate meter. A jugular vein was cannulated for the administration of drug and allergen. After surgery the animals were allowed to stabilise and the drug was administered i.v. as a bolus. Following this, ovalbumen 1mg.kg<sup>-1</sup> was injected i.v. as the antigen challenge either 2, 15 or 30 minutes following drug treatment and the peak bronchoconstrictor response recorded. For the control group ovalbumen only was given. One ovalbumen challenge per guinea-pig was used and  $n = 6$  for each time point. The percentage increase in tracheal inflation pressure was calculated. The following results indicating an anti-allergic activity were obtained.

Compound of Example	Dose $\mu$ mol/kg	% Inhibition of Control Bronchoconstrictor Response 30 min after drug administration
3	18.4	24
15	16.6	30

#### Phosphodiesterase activity

The activity of the compounds of the present invention as inhibitors of a calmodulin insensitive cyclic GMP phosphodiesterase was measured using the procedure described in European Patent Application No.

293063. The compounds of Examples 2-4, 6, 8, 9 and 11 to 15 had  $IC_{50}$  values (the concentration of inhibitor required for 50% inhibition of enzyme activity) in the range 0.31 to 4.80  $\mu$ M. The compounds of the present invention have the advantage that they are selective in not inhibiting cyclic AMP phosphodiesterase (type III).

#### Description 1

##### 2-(2-Propoxyphenyl)-6-purinone

A stirred mixture of 4,5-diamino-2-(2-propoxyphenyl)-pyrimidin-6-one sulphate (1.5 g) (prepared by the addition of concentrated sulphuric acid to an ethanolic solution of the free base) and formamide (15 ml) was heated in an oil bath (temp. 190°-200°C) for 70 minutes. When cool the mixture was filtered and the collected solid was washed with ethanol to give a crude product (1.1 g), m.p. 254-259°C, which was recrystallised from ethanol to give the title compound, 0.72 g, m.p. 263-265°C.

#### Description 2

##### 2-(2-Propoxyphenyl)purine-6,8-dione

A mixture of 4,5-diamino-2-(2-propoxyphenyl)pyrimidin-6-one (1.3 g), and urea (1.5 g) was heated in an oil bath (temp. 190°C) for 45 minutes. The resultant solid was digested with hot water, the mixture filtered and the solid washed with water to give a crude product, 1.36 g. Recrystallisation from dimethylformamide gave the title compound (1.01 g), m.p. >350°C,  $\delta$ (DMSO- $d_6$ ), 1.01 (t, 3H); 1.88 (m, 2H); 4.09 (t, 2H); 7.10, 7.21, 7.52 and 7.76 (multiplets, 4H);  $\tau$  11.07, 11.55 and 11.95 (very broad singlets, 3H).

#### Example 1

##### 2-(2-[2,2,2-Trifluoroethoxy]phenyl)purin-6-one

- a) A solution of 2-(2,2,2-trifluoromethoxy)benzamide (17 g, known from US Patent 3,766,247) and triethyloxonium tetrafluoroborate (ca. 28 g) in dichloromethane (140 ml) was allowed to stand for 20 hours. The solution was washed with saturated sodium carbonate solution, brine, dried over magnesium sulphate and evaporated to low volume under reduced pressure. The addition of ether (100 ml) and concentrated hydrochloric acid (6.5 ml) gave a solid which was recrystallised from ethanol-ether to give ethyl 2-(2,2,2-trifluoromethoxy)benzimidate hydrochloride, 12.88 g, m.p. 142.5-144°C (after transition 119-121°C).
- b) A solution of the above benzimidate hydrochloride (12.6 g) in saturated methanolic ammonia (75 ml) was allowed to stand for 40 hours. Evaporation to low volume gave a slurry which was diluted with ether to give 2-(2,2,2-trifluoromethoxy)benzamidine hydrochloride, 10.84 g, m.p. 248-251°C. Recrystallisation from ethanol-ether gave an analytical sample m.p. 250-252°C.
- c) A stirred mixture of the above benzamidine hydrochloride (10 g) and ethyl cyanoglyoxylate oxime (7.2 g) in sodium ethoxide solution (from sodium, 3.6 g, and ethanol, 300 ml) was heated under reflux for 5 hours. The mixture was evaporated under reduced pressure to a quarter volume, diluted with cold water (400 ml) and 2 Normal hydrochloric acid was added to pH 6. Filtration gave 4-amino-5-nitroso-2-(2-[2,2,2-trifluoroethoxy]phenyl)pyrimidin-6-one, 5.79 g, m.p. 210-212°C, which was used directly in the next stage.
- d) Sodium dithionite (3.88 g) was added during 5 minutes to a stirred partial solution of the above nitroso compound (3.5 g) and sodium biocarbonate (0.95 g) in 50% aqueous acetonitrile (180 ml) at 65°C. The resultant solution was stirred at 70°C for a further 10 minutes and then the bulk of the acetonitrile was removed by evaporation under reduced pressure. The cold mixture was filtered to give crude 4,5-diamino-2-(2-[2,2,2-trifluoroethoxy]phenyl)pyrimidin-6-one, which was dissolved in the minimum volume of hot ethanol and converted into the sulphate salt (2.33 g, m.p. 255-260°C dec) by the addition of sulphuric acid. Recrystallisation from 50% aqueous ethanol gave the hemisulphate as a partial hydrate, m.p. ca. 210-220°C dec.
- e) A stirred mixture of the above diamine hemisulphate (1.1 g) and formamide (11 ml) was heated in an oil bath (temperature 195°C) for 2 hours. The solution was cooled and diluted with water (44 ml) to give 0.92 g of a solid m.p. 261-263°C. Recrystallisation from ethanol gave the pure title compound, 0.77 g, m.p. 276-277°C (transition 263°C).

Example 22-(2-Cyclopropylmethoxyphenyl)purin-6-one

5 A stirred mixture of 4,5-diamino-2-(2-cyclopropylmethoxyphenyl)pyrimidin-6-one sulphate (1.48 g) and formamide (5 ml) was heated in an oil bath at 180 °C for 2 to 3 hours. The cooled mixture was filtered and the collected solid was washed with ethanol to give a crude product (1.07 g) which was recrystallised three times from ethanol to afford the title compound, 0.32 g, m.p. 259-260 °C.

Example 32-(2-Cyclopropylmethoxyphenyl)purin-6,8-dione

15 A mixture of 4,5-diamino-2-(2-cyclopropylmethoxyphenyl)pyrimidin-6-one (0.90 g) and urea (0.99 g) was heated in an oil bath at 160 ° to 170 °C for one hour. The resultant solid was digested with warm water and the mixture filtered to afford a solid (0.74 g) which was twice recrystallised from dimethylformamide to afford a crude product (0.34 g). This together with another sample (0.22 g) similarly prepared from 4,5-diamino-2-(2-cyclopropylmethoxyphenyl)pyrimidin-6-one (0.45 g) was twice recrystallised from dimethylformamide to afford the title compound, 0.34 g, m.p. 329-331 °C.

Example 42-(2-Benzoyloxyphenyl)purin-6,8-dione

25 2-(2-Benzoyloxyphenyl)-4-amino-5-nitrosopyrimidine-6-one (2.0 g) was suspended in 50 ml of 1:1 acetonitrile : water and heated to 70 °C. A solution of sodium dithionite (1.9 g) in water (10 ml) was added dropwise over 5 minutes and heating continued for a further 10 minutes. The solution was cooled to room temperature, poured into saturated aqueous sodium hydrogen carbonate (250 ml) and extracted with dichloromethane. The organic extract was dried (magnesium sulphate), concentrated to ca 50 ml and treated with carbonyl diimidazole (1.3 g). After 16 hours, solvents were removed in vacuo and the residue recrystallised from dimethylformamide/water to afford the title compound, 1.2 g, m.p. 295 °C (dec).

Example 52-(2-propoxyphenyl)-8-trifluoromethylpurin-6-one

35 a) 4,5-Diamino-2-(2-propoxyphenyl)pyrimidin-6-one sulphate (1 g) and trifluoroacetic anhydride (10 ml) were heated together under reflux for 2 hours. Potassium carbonate (0.38 g) was added and the mixture was heated for a further 2 hours. The residue left after evaporation was treated with water (25 ml) and potassium carbonate was added to pH 5 to give 0.92 g of solid m.p. 192-199 °C. Purification by column chromatography (silica gel, chloroform) gave 0.88 g of a solid m.p. 198-201 °C, which was recrystallised from isopropyl acetate and then acetonitrile to give an analytical sample of 5-trifluoroacetyl-amino-4-amino-2-(2-propoxyphenyl)pyrimidin-6-one, m.p. 203-204 °C.

45 b) A melt of the above trifluoroacetyl-amino derivative (0.62 g) was heated under nitrogen in an oil bath (temperature 250 °C) for 10 minutes to give a solid, m.p. 255-260 °C. Recrystallisation from ethanol gave the pure title compound, 0.3 g, m.p. 267-269 °C.

Example 62-(2-Propoxyphenyl)-8-phenylpurin-6-one

50 A mixture of 4,5-diamino-2-(2-propoxyphenyl)-pyrimidine-6-one (1.3 g), benzamidine hydrochloride (2 g) and anhydrous sodium acetate (0.9 g) was heated in an oil bath (temperature 160-170 °C) for 2 hours. The reaction mixture was digested with hot ethanol to give a solid, 0.76 g, m.p. 267-271 °C. This was combined with a further sample, 0.2 g, (similarly prepared) and recrystallised from aqueous acetic acid to give the title compound, 0.74 g, m.p. 259-260 °C.



Example 72-(2-Propoxyphenyl)-8-methylpurin-6-one

5 A mixture of 4,5-diamino-2-(2-propoxyphenyl)pyrimidin-6-one (1.56 g), anhydrous sodium acetate (1.14 g) and acetamidine hydrochloride (1.42 g) was heated in an oil bath at 150-160 °C for 2 hours. The mixture was digested with ethanol (2 ml), cooled, filtered and the solid washed with ethanol. Recrystallisation from ethanol gave the title compound, 0.72 g, m.p. 264-265 °C.

Example 82-(2-Propoxyphenyl)-8-mercaptopurin-6-one

15 A mixture of 4,5-diamino-2-(2-propoxyphenyl)pyrimidin-6-one (1.3 g), anhydrous potassium acetate (0.34 g) and thiourea (0.96 g) was heated at 170-180 °C for 2 hours. After digestion with water the mixture was filtered and the solid was dissolved in 1 Normal sodium hydroxide solution then re-precipitated by the addition of acetic acid. The dried solid was digested with chloroform and with ethanol to leave 1.12 g of a crude product. Repeated recrystallisations from dimethylformamide, 2-methoxyethanol, and acetic acid yielded the title compound, 0.32 g, m.p. 305-307 °C.

20

Example 92-(2-Propoxyphenyl)-8-methylthiopurin-6-one

25 Methyl iodide (0.5 g) was added to a stirred solution of 2-(2-propoxyphenyl)-8-mercaptopurin-6-one (1.0 g) in 1 Normal sodium hydroxide solution (8 ml). After 2.5 hours at room temperature the solution was neutralised with dilute hydrochloric acid to give the crude product which was recrystallised three times from aqueous ethanol to give the title compound, 0.32 g, m.p. 245-247 °C.

Example 102-(2-Propoxyphenyl)-8-aminopurin-6-one

35 A partial solution of 4,5-diamino-2-(2-propoxyphenyl)pyrimidin-6-one sulphate (1.1 g), cyanogen bromide (0.33 g), and sodium acetate trihydrate (0.42 g) in 50% aqueous ethanol (44 ml) was stirred at room temperature for 2 hours then allowed to stand overnight. The stirred mixture was then heated in a water bath (temperature 65 °C) for 3 hours, extra cyanogen bromide (0.05 g) added, and the mixture heated for a further 2 hours. The suspension was subjected to partial evaporation under reduced pressure then ammonium hydroxide was added to pH 5. The crude product was collected by filtration and recrystallised  
40 from acetonitrile to give the pure title compound, 0.6 g, m.p. 322-335 °C dec. (after melting ca. 200 °C).  
NMR (DMSO-d<sub>6</sub>; 250MHz) δ : 0.9 (3H, t); 1.6 (2H, m); 3.9 (2H, t); 6.8 (2H, s); 6.8 (2H, s); 6.9-7.1 (2H, m); 7.3-7.5 (3H, m).

Example 11

45

2-(2-Propoxy-5-nitrophenyl)purin-6-one

A mixture of fuming nitric acid (0.23 ml) and sulphuric acid (4 ml) was added dropwise to a stirred solution of 2-(2-propoxyphenyl)purin-6-one (1.0 g) in sulphuric acid (4 ml) at 0 to -5 °C. The temperature  
50 was maintained between -5 °C and +4 °C for 20 hours and then the mixture was poured into ice-water. The filtered solution was treated with concentrated ammonium hydroxide to pH 9 to give a crude product, 0.55 g. Recrystallisation twice from aqueous ethanol then once from acetonitrile gave the title compound, 0.2 g, m.p. 254-256 °C.

55

Example 122-(2-Propoxy-5-acetamidophenyl)purin-6-one

5 A solution of the crude product of Example 11 (1.5 g) in water (50 ml) containing 2 Normal sodium hydroxide (2.3 ml) and 10% palladium on charcoal (0.15 g) was shaken under hydrogen (50 psi) until the uptake was complete. Neutralisation of the filtered solution with acetic acid gave a fine precipitate of 2-(5-amino-2-propoxyphenyl)-purin-6-one. This mixture was warmed with 2 Normal hydrochloric acid (2.5 ml) and the solution was treated with acetic anhydride (0.55 ml) and sodium acetate trihydrate (0.8 g). The  
 10 mixture was warmed for 10 minutes then cooled and filtered to give a crude product (1.02 g) which was recrystallised from aqueous dimethylformamide twice to give the title compound, 0.49 g, m.p. 320-323 °C.

Example 132-(2-propoxy-4-methoxyphenyl)purin-6-one

a) A stirred mixture of methyl 4-methoxysalicylate (25 g), bromopropane (15.6 ml), potassium iodide (2.82 g) and anhydrous potassium carbonate (27.53 g) was heated under reflux for 48 hours. The cooled reaction mixture was filtered and the filtrate was evaporated under reduced pressure to yield an oil which  
 20 was dissolved in diethyl ether (200 ml). The ethereal solution was extracted with aqueous sodium hydroxide to remove unreacted starting material and the organic phase was then washed with water and brine, dried (magnesium sulphate) and evaporated under reduced pressure to yield methyl 4-methoxy-2-propoxy-benzoate, 22.45 g.

b) Methyl 4-methoxy-2-propoxybenzoate, (22.35 g) was treated with a saturated solution of ammonia in dry methanol (150 ml) for 6 hours at 80 °C in a pressure vessel. From the cooled reaction mixture was  
 25 collected as a precipitate a crude sample of 4-methoxy-2-propoxybenzamide, 8.9 g. Recrystallisation from acetonitrile gave an analytical sample, m.p. 130-132 °C.

c) A mixture of 4-methoxy-2-propoxybenzamide (15 g) and triethyloxonium tetrafluoroborate (0.08 mol) in dichloromethane (180 ml) was stirred at ambient temperature for about 60 hours. The reaction mixture  
 30 was evaporated under reduced pressure and the residue was washed with diethyl ether to yield crude ethyl 4-methoxy-2-propoxybenzimidate tetrafluoroborate, 20.52 g which was used without further purification.

d) A mixture of the above imidate salt (20.40 g) and saturated ethanolic ammonia (150 ml) was stirred for 18 hours at ambient temperature. Excess ammonia was removed by evaporation on a steam bath and  
 35 the reaction mixture was evaporated under reduced pressure to low volume (50 ml). Concentrated hydrochloric acid (8 ml) was added and the mixture was evaporated under reduced pressure to yield a residue which was triturated with diethyl ether and a little ethanol to yield 4-methoxy-2-propoxybenzimidine hydrochloride, 7.67 g.

e) A stirred mixture of the above benzimidine hydrochloride (7.63 g) and ethyl cyanoglyoxylate-2-oxime (4.43 g) in sodium ethoxide solution (from sodium, 2.85 g, and ethanol, 120 ml) was heated under reflux for 3.5 hours. The cooled reaction mixture was evaporated to dryness and the residue was dissolved in  
 40 water. Addition of hydrochloric acid yielded a precipitate which was collected, washed with water, digested with warm ethanol and finally washed with ethanol and diethyl ether to yield a solid (2.50 g). This was stirred in dilute hydrochloric acid for 10 minutes, filtered and washed with water to yield 2-(4-methoxy-2-propoxyphenyl)-4-amino-5-nitroso-pyrimidin-6-one, 2.16 g, m.p. 242-245 °C (dec).

f) A stirred mixture of the above nitroso compound (2.10 g), sodium bicarbonate (1.28 g) and sodium dithionite (2.64 g) in 50% aqueous acetonitrile (150 ml) was heated at 70 °C for 10 minutes and then chilled for 30 minutes. A two phase system formed. The upper organic phase was separated, washed  
 50 with brine and reduced in volume to about 5 ml. This was dissolved in ethanol (20 ml), treated with concentrated sulphuric acid (1 ml) and concentrated by evaporation until precipitation began to occur. The cooled mixture yielded 2-(4-methoxy-2-propoxyphenyl)-4,5-diaminopyrimidin-6-one sulphate, 1.06 g, m.p. 209-212 °C (dec). The aqueous phase was concentrated to about half volume and was extracted with chloroform (3x 25 ml). The combined organic extracts were washed with water and brine, dried (magnesium sulphate) and evaporated to dryness. The residue was dissolved in ethanol (15 ml) and  
 55 treated with concentrated sulphuric acid (0.5 ml) and the resulting solution was evaporated until precipitation occurred. A little diethyl ether was added and the cooled mixture afforded a further sample of the above diamine sulphate, 0.26 g, m.p. 213-6 °C (dec).

g) A stirred mixture of the above diamine sulphate (0.75 g) and formic acid (5 ml) was heated under reflux for 4.5 hours. The cooled reaction mixture was poured into water (25 ml) and the resultant mixture was centrifuged for 15 minutes to yield a solid (0.60 g) which together with another sample (0.25 g), similarly prepared, was recrystallised from 50% aqueous ethanol to afford the title compound, 0.60 g, m.p. 290-1 °C (dec).

#### Example 14

##### 2-(2-Propoxy-5-methoxyphenyl)purin-6-one

In a similar manner to Example 13 :

a) reaction of 5-methoxy-2-propoxybenzamide (10.88 g) with triethyloxonium tetrafluoroborate (0.07 mol) yielded ethyl 5-methoxy-2-propoxybenzimidate tetrafluoroborate (21.0 g);

b) reaction of the above imidate salt (16.92 g) with a saturated solution of ethanolic ammonia (150 ml) yielded crude 5-methoxy-2-propoxybenzamidine (9.98 g);

c) reaction of the above amidine (9.98 g) with ethyl cyanoglyoxylate-2-oxime (6.86 g) and sodium ethoxide (from sodium, 3.31 g, and ethanol, 100 ml) yielded 2-(5-methoxy-2-propoxyphenyl)-4-amino-5-nitrosopyrimidin-6-one (6.63 g);

d) reaction of the above nitroso compound (3.80 g) with sodium dithionite (4.83 g) and sodium bicarbonate (2.33 g) yielded on treatment with concentrated sulphuric acid 2-(5-methoxy-2-propoxyphenyl)-4,5-diaminopyrimidin-6-one sulphate, 2.49 g, m.p. 221-224 °C (dec);

e) reaction of the above diamine sulphate (1.80 g) with formic acid (10 ml) yielded the title compound, 0.61 g, m.p. 233-4 °C (recrystallised from 25% aqueous ethanol).

#### Example 15

##### 2-(2-Propoxy-5-chlorophenyl)purin-6-one

In a similar manner to Example 13 :

a) reaction of 5-chloro-2-propoxybenzamide (16.50 g) with triethyloxonium tetrafluoroborate (0.096 mol) yielded crude ethyl 5-chloro-2-propoxybenzimidate tetrafluoroborate (29.14 g);

b) reaction of the above imidate salt (29.14 g) with a saturated solution of ethanolic ammonia (200 ml) yielded 5-chloro-2-propoxybenzamidine (9.30 g);

c) reaction of the above amidine (4.50 g) with ethyl cyanoglyoxylate-2-oxime (4.43 g) and sodium ethoxide (from sodium, 1.46 g, and ethanol, 100 ml) yielded 2-(5-chloro-2-propoxyphenyl)-4-amino-5-nitrosopyrimidin-6-one (1.89 g);

d) reaction of the above nitroso compound (1.02 g) with sodium dithionite (1.26 g) and sodium bicarbonate (0.61 g) yielded on treatment with concentrated sulphuric acid 2-(5-chloro-2-propoxyphenyl)-4,5-diaminopyrimidin-6-one sulphate, 0.81 g, m.p. 220-223 °C;

e) reaction of the above diamine sulphate (0.60 g) with formic acid (3 ml) yielded the crude title compound, (0.48 g) which together with another sample (0.48 g), similarly prepared, was recrystallised from 50% aqueous ethanol to afford the title compound, 0.61 g, m.p. 277-9 °C.

The starting-material, 5-chloro-2-propoxybenzamide, was prepared as follows :

A stirred mixture of 5-chloro-2-hydroxybenzamide (20 g), 1-bromopropane (13.4 ml), potassium iodide (2.49 g) and anhydrous potassium carbonate (24.15 g) in acetone (250 ml) was heated under reflux for 20 hours. The cooled reaction mixture was filtered and the filter cake was washed with acetone. The filtrate and washings were combined and evaporated under reduced pressure to yield a residue which was washed with water, dilute aqueous sodium hydroxide, water and with diethyl ether to yield a crude product (19.85 g). This was recrystallised from acetonitrile, partitioned between chloroform and dilute aqueous sodium hydroxide, and was recrystallised from ethanol to yield 5-chloro-2-propoxybenzamide (16.78 g).

#### Example 16

##### 2-(2-Propoxy-4-methylphenyl)purin-6-one

In a similar manner to Example 13 :

a) reaction of 4-methyl-2-propoxybenzamide (11.96 g) with triethyloxonium tetrafluoroborate yielded crude ethyl 4-methyl-2-propoxybenzimidate tetrafluoroborate which on reaction with a saturated solution

of ethanolic ammonia and treatment with hydrochloric acid yielded 4-methyl-2-propoxybenzamidine hydrochloride, (8.74 g), m.p. 228-30 °C;

b) reaction of the above amidine (8.70 g) with ethyl cyanoglyoxylate-2-oxime (4.80 g) and sodium ethoxide yielded 2-(4-methyl-2-propoxyphenyl)-4-amino-5-nitrosopyrimidin-6-one, 3.26 g, m.p. 214-6 °C;

5 c) reaction of the above nitroso compound (3.24 g) with sodium dithionite and sodium bicarbonate yielded on treatment with concentrated sulphuric acid 2-(4-methyl-2-propoxyphenyl)-4,5-diaminopyrimidin-6-one sulphate, 4.24 g;

d) reaction of the above diamine sulphate (4.24 g) with formic acid yielded the crude title compound, which was recrystallised from aqueous dimethylformamide to yield the title compound, 1.09 g, m.p. 323-5 °C.

The starting-material, 4-methyl-2-propoxybenzamide, was prepared by reacting methyl 4-methylsalicylate (25.75 g) with ethanolic ammonia to yield 4-methylsalicylamide (12.54 g) which was then reacted with bromopropane, potassium iodide and potassium carbonate in acetone.

#### 15 Example 17

##### 2-(2-Propoxy-5-fluorophenyl)purin-6-one

In a similar manner to Example 13 :

20 a) reaction of 5-fluoro-2-propoxybenzamide (16.40 g) with triethyloxonium tetrafluoroborate yielded ethyl 5-fluoro-2-propoxybenzimidate tetrafluoroborate which on reaction with a saturated solution of ethanolic ammonia yielded 5-fluoro-2-propoxybenzamidine (8.40 g);

b) reaction of the above amidine (5.0 g) with ethyl cyanoglyoxylate-2-oxime (3.35 g) and sodium methoxide (from sodium, 2.17 g, and methanol, 200 ml) yielded 2-(5-fluoro-2-propoxyphenyl)-4-amino-5-nitrosopyrimidin-6-one, 2.2 g, m.p. 220 °C (dec);

25 c) reaction of the above nitroso compound (1.0 g) with sodium dithionite (1.1 g) yielded 2-(5-fluoro-2-propoxyphenyl)-4,5-diaminopyrimidin-6-one, 0.75 g, which on reaction with formic acid (10 ml) yielded the crude title compound, (0.63 g) which was recrystallised from aqueous dimethylformamide to yield the title compound, 0.40 g, m.p. 238 °C.

30 The starting-material, 5-fluoro-2-propoxybenzamide, was prepared as follows :

Alkylation of 5-fluoro-2-hydroxyacetophenone (25 g) with bromopropane, potassium iodide and potassium carbonate in acetone yielded 5-fluoro-2-propoxyacetophenone, 20.9 g, m.p. 59-61 °C. 5-Fluoro-2-propoxyacetophenone, (1 g) was shaken in a solution of bromine (2 ml) in aqueous sodium hydroxide (3.6 g in 30 ml) for 20 minutes, then the stirred mixture was heated at 90 °C for 4 hours. Excess of hypobromite was destroyed with sodium metabisulphite, and the solution was extracted with ether. The organic solution was extracted with 2 Normal sodium hydroxide, the extract acidified and extracted with dichloromethane. Evaporation yielded 5-fluoro-2-propoxybenzoic acid, 0.32 g, m.p. 76-78 °C. 5-Fluoro-2-propoxybenzoic acid (29 g) was treated with thionyl chloride to yield the corresponding acid chloride, which was in turn treated with ammonia in ether solution to give 5-fluoro-2-propoxybenzamide, 16.5 g, m.p. 114-116 °C.

#### 40 Example 18

##### 2-(2-Propoxy-5-dimethylsulphamoylphenyl)purin-6-one

45 2-(2-Propoxyphenyl)purin-6-one (0.27 g) was added portionwise with cooling (0 °C) to stirred chlorosulphonic acid (1.25 ml). The reaction mixture was stirred with cooling (0 °C) for 20 minutes and then left at 4 °C for 24 hours. The mixture was washed with dichloromethane (2 x 20 ml) and the residue was added with cooling (5 °C) to a stirred solution of dimethylamine in industrial methylated spirit (33%, 10 ml). The resulting solution was stirred at ambient temperature for 30 minutes and evaporated under reduced pressure to yield an oil which was treated with water (15 ml) and made basic to pH9 with potassium carbonate. After standing overnight at ambient temperature a precipitate (0.18 g) was collected, which together with another sample (0.50 g), similarly prepared from 2-(2-propoxyphenyl)purin-6-one (0.68 g) and chlorosulphonic acid (3.5 ml), was recrystallised from aqueous ethanol (with charcoal) to yield the title compound, 0.43 g, m.p. 268-9 °C.

55

Example 192-(2-Propoxy-5-methylsulphamoylphenyl)purin-6-one

5 In a similar manner to Example 18 reaction of 2-(2-propoxyphenyl)purin-6-one (0.81 g) with chlorosulphonic acid (4.5 ml), followed by reaction with methylamine in industrial methylated spirit (33%, 50 ml) yielded the title compound, 0.49 g, m.p. 245-246 °C (recrystallised twice from aqueous ethanol).

Example 202-(2-Propoxy-5-sulphamoylphenyl)purin-6-one

10 In a similar manner to Example 18 reaction of 2-(2-propoxyphenyl)purin-6-one (0.95 g) with chlorosulphonic acid (4.5 ml), followed by reaction with saturated methanolic ammonia (50 ml), yielded a crude product (0.92 g) which was recrystallised from aqueous ethanol and then from dimethylformamide to yield the pure title compound, 0.59 g, m.p. 276-278 °C.

Example 21

20 Pharmaceutical compositions for oral administration are prepared by combining the following :

	% w/w		
25 2-(2-Propoxy-5-chlorophenyl)purin-6-one	0.5	3.0	7.14
2% w/w Soya lecithin in soya bean oil	90.45	88.2	84.41
Hydrogenated vegetable shortening and beeswax	9.05	8.8	8.45

The formulations are then filled into individual soft gelatin capsules.


Example 22

30 A pharmaceutical composition for parenteral administration is prepared by dissolving the title compound of Example 3 (0.02 g) in polyethylene glycol 300 (25 ml) with heating. This solution is then diluted with water for injections Ph. Eur. (to 100 ml). The solution is then sterilised by filtration through a 0.22 micron membrane filter and sealed in sterile containers.

Claims

Claims for the following Contracting States : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

40 1. A compound of the formula (1) :



or a pharmaceutically acceptable salt thereof, wherein

R¹ is C₁-₆ alkyl, C₂-₆ alkenyl, C₃-₅ cycloalkyl, C₁-₄ alkyl, phenyl, C₁-₄ alkyl or C₁-₄ alkyl substituted by 1 to 6 fluoro groups;

55 R² is hydrogen, hydroxy, C₁-₄ alkyl, phenyl, mercapto, C₁-₄ alkylthio, CF₃ or amino;


R³ is hydrogen, nitro, amino, C₁-₄ alkanoylamino, C₁-₄ alkoxy, C₁-₄ alkyl, halo, SO₂NR⁴R⁵, CONR⁴R⁵, cyano or C₁-₄ alkylS(O)ₙ;

R⁴ and R⁵ are independently hydrogen or C₁-₄ alkyl; and

n is 0, 1 or 2;  
provided that R<sup>3</sup> is not hydrogen when R<sup>1</sup> is C<sub>1-6</sub> alkyl or C<sub>2-6</sub> alkenyl and R<sup>2</sup> is hydrogen or hydroxy.

2. A compound according to claim 1 wherein R<sup>1</sup> is C<sub>2-5</sub> alkyl.
3. A compound according to claim 1 wherein R<sup>1</sup> is C<sub>3-5</sub> alkenyl.
4. A compound according to claim 1 wherein R<sup>1</sup> is cyclopropylmethyl.
5. A compound according to claim 1 wherein R<sup>1</sup> is CF<sub>3</sub>, CH<sub>2</sub>CF<sub>3</sub> or CF<sub>2</sub>CHFCH<sub>2</sub>CF<sub>3</sub>.
6. A compound according to any one of claims 1 to 5 wherein R<sup>2</sup> is hydrogen or hydroxy.
7. A compound according to any one of claims 1 to 5 wherein R<sup>2</sup> is phenyl or C<sub>1-4</sub> alkyl.
8. A compound according to any one of claims 1 to 5 wherein R<sup>2</sup> is mercapto or C<sub>1-4</sub> alkylthio.
9. A compound according to any one of claims 1 to 8 wherein R<sup>3</sup> is hydrogen.
10. A compound according to any one of claims 1 to 8 wherein R<sup>3</sup> is nitro, cyano, CONR<sup>4</sup>R<sup>5</sup> or SO<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>.
11. A compound according to any one of claims 1 to 8 wherein R<sup>3</sup> is halo, C<sub>1-4</sub> alkanoylamino, C<sub>1-4</sub> alkoxy or C<sub>1-4</sub> alkyl.
12. A compound according to claim 1 which is :  
 2-(2-[2,2,2-trifluoroethoxy]phenyl)purin-6-one,  
 2-(2-cyclopropylmethoxyphenyl)purin-6-one,  
 2-(2-cyclopropylmethoxyphenyl)purin-6,8-dione,  
 2-(2-benzoyloxyphenyl)purin-6,8-dione,  
 2-(2-propoxyphenyl)-8-trifluoromethylpurin-6-one,  
 2-(2-propoxyphenyl)-8-phenylpurin-6-one,  
 2-(2-propoxyphenyl)-8-methylpurin-6-one,  
 2-(2-propoxyphenyl)-8-mercaptapurin-6-one,  
 2-(2-propoxyphenyl)-8-methylthiopurin-6-one,  
 2-(2-propoxyphenyl)-8-aminopurin-6-one,  
 2-(2-propoxy-5-nitrophenyl)purin-6-one,  
 2-(2-propoxy-5-aminophenyl)purin-6-one,  
 2-(2-propoxy-5-acetamidophenyl)purin-6-one,  
 2-(2-propoxy-4-methoxyphenyl)purin-6-one,  
 2-(2-propoxy-5-methoxyphenyl)purin-6-one,  
 2-(2-propoxy-5-chlorophenyl)purin-6-one,  
 2-(2-propoxy-4-methylphenyl)purin-6-one,  
 2-(2-propoxy-5-fluorophenyl)purin-6-one,  
 2-(2-propoxy-5-dimethylsulphamoylphenyl)purin-6-one,  
 2-(2-propoxy-5-methylsulphamoylphenyl)purin-6-one,  
 2-(2-propoxy-5-sulphamoylphenyl)purin-6-one,  
 2-(2-propoxy-4-methylthiophenyl)purin-6-one,  
 2-(2-propoxy-5-cyanophenyl)purin-6-one, or  
 2-(2-propoxy-5-carbamoylphenyl)purin-6-one,  
 or a pharmaceutically acceptable salt thereof.
13. A compound according to any one of claims 1 to 12 for use as a medicament.
14. A pharmaceutical composition which comprises a compound according to any one of claims 1 to 12 and a pharmaceutically acceptable carrier.
15. A process for preparing a compound of the formula (1) or a pharmaceutically acceptable salt thereof as defined in claim 1 which comprises :

a) for compounds wherein  $R^2$  is hydrogen,  $C_1-4$ -alkyl, phenyl or  $CF_3$ ,  
reacting a compound of the formula (2) :



15 wherein  $R^1$  as defined in claim 1 and  $R^6$  is a group  $R^3$  as defined in claim 1 or a precursor thereof,  
with a compound of the formula  $R^7COL$  or a chemical equivalent thereof wherein  $R^7$  is hydrogen,  
 $C_1-4$ alkyl, phenyl or  $CF_3$  and  $L$  is a leaving group;

b) for compounds wherein  $R^2$  is hydroxy,

reacting a compound of the formula (2) as hereinbefore defined with a carbonylating agent;

c) for compounds wherein  $R^2$  is mercapto,

20 reacting a compound of the formula (2) as hereinbefore defined with thiourea;

d) for compounds wherein  $R^2$  is  $C_1-4$ alkylthio,

reacting a compound of the formula (2) as hereinbefore defined with thiourea and thereafter with a  
 $C_1-4$ alkyl halide;

e) for compounds wherein  $R^2$  is amino,


25 reacting a compound of the formula (2) as hereinbefore defined with cyanogen bromide;

and thereafter where necessary :

- converting a group  $R^6$  to a group  $R^3$ ;
- optionally forming a pharmaceutically acceptable salt.

### 30 Claims for the following Contracting States: ES, GR

1. A process for preparing a compound of the formula (1) :



or a pharmaceutically acceptable salt thereof, wherein

45  $R^1$  is  $C_1-6$ alkyl,  $C_2-6$ alkenyl,  $C_3-5$ cycloalkyl,  $C_1-4$ alkyl, phenyl,  $C_1-4$ alkyl or  $C_1-4$ alkyl  
substituted by 1 to 6 fluoro groups;

$R^2$  is hydrogen, hydroxy,  $C_1-4$ alkyl, phenyl, mercapto,  $C_1-4$ alkylthio,  $CF_3$  or amino;


$R^3$  is hydrogen, nitro, amino,  $C_1-4$ alkanoylamino,  $C_1-4$ -alkoxy,  $C_1-4$ alkyl, halo,  
50  $SO_2NR^4R^5$ ,  $CONR^4R^5$ , cyano or  $C_1-4$ alkylS(O) $_n$ ;

$R^4$  and  $R^5$  are independently hydrogen or  $C_1-4$ alkyl; and

$n$  is 0, 1 or 2;

provided that  $R^3$  is not hydrogen when  $R^1$  is  $C_1-6$ alkyl or  $C_2-6$ alkenyl and  $R^2$  is hydrogen or hydroxy;  
which process comprises:

a) for compounds wherein  $R^2$  is hydrogen,  $C_1-4$ alkyl, phenyl or  $CF_3$ , reacting a compound of the  
55 formula (2):



(2)

wherein R<sup>1</sup> is as hereinbefore defined and R<sup>6</sup> is a group R<sup>3</sup> as hereinbefore defined or a precursor thereof, with a compound of the formula R<sup>7</sup>COL or a chemical equivalent thereof wherein R<sup>7</sup> is hydrogen, C<sub>1-4</sub>alkyl, phenyl or CF<sub>3</sub> and L is a leaving group;

b) for compounds wherein R<sup>2</sup> is hydroxy,

reacting a compound of the formula (2) as hereinbefore defined with a carbonylating agent;

c) for compounds wherein R<sup>2</sup> is mercapto,

reacting a compound of the formula (2) as hereinbefore defined with thiourea;

d) for compounds wherein R<sup>2</sup> is C<sub>1-4</sub>alkylthio,

reacting a compound of the formula (2) as hereinbefore defined with thiourea and thereafter with a C<sub>1-4</sub>alkyl halide;

e) for compounds wherein R<sup>2</sup> is amino, reacting a compound of the formula (2) as hereinbefore defined with cyanogen bromide;

and thereafter where necessary :

- converting a group R<sup>6</sup> to a group R<sup>3</sup>;

- optionally forming a pharmaceutically acceptable salt.

2. A process according to claim 1 for preparing a compound wherein R<sup>1</sup> is C<sub>2-5</sub>alkyl.

3. A process according to claim 1 for preparing a compound wherein R<sup>1</sup> is C<sub>3-5</sub>alkenyl.

4. A process according to claim 1 for preparing a compound wherein R<sup>1</sup> is cyclopropylmethyl.

5. A process according to claim 1 for preparing a compound wherein R<sup>1</sup> is CF<sub>3</sub>, CH<sub>2</sub>CF<sub>3</sub> or CF<sub>2</sub>CHFCF<sub>3</sub>.

6. A process according to any one of claims 1 to 5 for preparing a compound wherein R<sup>2</sup> is hydrogen or hydroxy.

7. A process according to any one of claims 1 to 5 for preparing a compound wherein R<sup>2</sup> is phenyl or C<sub>1-4</sub>alkyl.

8. A process according to any one of claims 1 to 5 for preparing a compound wherein R<sup>2</sup> is mercapto or C<sub>1-4</sub>alkylthio.

9. A process according to any one of claims 1 to 8 for preparing a compound wherein R<sup>3</sup> is hydrogen.

10. A process according to any one of claims 1 to 8 for preparing a compound wherein R<sup>3</sup> is nitro, cyano, CONR<sup>4</sup>R<sup>5</sup> or SO<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>.

11. A process according to any one of claims 1 to 8 for preparing a compound wherein R<sup>3</sup> is halo, C<sub>1-4</sub>alkanoyl-amino, C<sub>1-4</sub>alkoxy or C<sub>1-4</sub>alkyl.

12. A process according to claim 1 for preparing a compound which is :

2-(2-[2,2,2-trifluoroethoxy]phenyl)purin-6-one,

2-(2-cyclopropylmethoxyphenyl)purin-6-one,

2-(2-cyclopropylmethoxyphenyl)purin-6,8-dione,

2-(2-benzoyloxyphenyl)purin-6,8-dione,

2-(2-propoxyphenyl)-8-trifluoromethylpurin-6-one,

2-(2-propoxyphenyl)-8-phenylpurin-6-one,




2-(2-propoxyphenyl)-8-methylpurin-6-one,  
 2-(2-propoxyphenyl)-8-mercaptapurin-6-one,  
 2-(2-propoxyphenyl)-8-methylthiopurin-6-one,  
 2-(2-propoxyphenyl)-8-aminopurin-6-one,  
 5 2-(2-propoxy-5-nitrophenyl)purin-6-one,  
 2-(2-propoxy-5-aminophenyl)purin-6-one,  
 2-(2-propoxy-5-acetamidophenyl)purin-6-one,  
 2-(2-propoxy-4-methoxyphenyl)purin-6-one,  
 2-(2-propoxy-5-methoxyphenyl)purin-6-one,  
 10 2-(2-propoxy-5-chlorophenyl)purin-6-one,  
 2-(2-propoxy-4-methylphenyl)purin-6-one,  
 2-(2-propoxy-5-fluorophenyl)purin-6-one,  
 2-(2-propoxy-5-dimethylsulphamoylphenyl)purin-6-one,  
 2-(2-propoxy-5-methylsulphamoylphenyl)purin-6-one,  
 15 2-(2-propoxy-5-sulphamoylphenyl)purin-6-one,  
 2-(2-propoxy-4-methylthiophenyl)purin-6-one,  
 2-(2-propoxy-5-cyanophenyl)purin-6-one, or  
 2-(2-propoxy-5-carbamoylphenyl)purin-6-one,  
 or a pharmaceutically acceptable salt thereof.

- 20 13. A process for preparing a pharmaceutical composition which comprises bringing into association a compound according to any one of claims 1 to 12 and a pharmaceutically acceptable carrier.
14. A process according to claim 1 wherein L in the compound of the formula  $R^7COL$  is selected from  
 25 hydroxy,  $C_1-4$ alkoxy, halo, amino,  $C_1-4$ alkylamino or  $OCOR^7$ .
15. A process according to claim 1 wherein a carbonylating agent is selected from urea,  $di(C_1-4)alkyl$  carbonate,  $C_1-4$ alkyl chloroformate, phosgene, trichloromethyl chloroformate or carbonyl diimidazole.

### 30 Patentansprüche

Patentansprüche für folgende Vertragsstaaten : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Verbindung der Formel (1):




(1),

45 oder ein pharmazeutisch verträgliches Salz davon, wobei  
 $R^1$  einen  $C_1-6$ -Alkyl-,  $C_2-6$ -Alkenyl-,  $C_3-5$ -Cycloalkyl- $C_1-4$ -alkyl-, Phenyl- $C_1-4$ -alkyl- oder einen mit 1 bis 6 Fluoratomen substituierten  $C_1-4$ -Alkylrest bedeutet,  
 $R^2$  ein Wasserstoffatom, eine Hydroxygruppe, einen  $C_1-4$ -Alkylrest, eine Phenyl- oder Mercaptogruppe,  
 50 einen  $C_1-4$ -Alkylthioester, eine Gruppe  $CF_3$  oder eine Aminogruppe bedeutet,  
 $R^3$  ein Wasserstoffatom, eine Nitro- oder Aminogruppe, einen  $C_1-4$ -Alkanoylamino-,  $C_1-4$ -Alkoxy- oder  
 $C_1-4$ -Alkylrest, ein Halogenatom, einen Rest  
 $-SO_2NR^4R^5$  oder  $-CONR^4R^5$ , eine Cyanogruppe oder einen Rest  $-C_1-4-AlkylS(O)_n$  bedeutet,  
 $R^4$  und  $R^5$  unabhängig voneinander ein Wasserstoffatom oder einen  $C_1-4$ -Alkylrest bedeuten, und  
 55  $n$  0, 1 oder 2 ist,  
 mit der Maßgabe, daß  $R^3$  kein Wasserstoffatom ist, wenn  $R^1$  einen  $C_1-6$ -Alkyl- oder  $C_2-6$ -Alkenylrest  
 bedeutet und  $R^2$  ein Wasserstoffatom oder eine Hydroxygruppe bedeutet.

2. Verbindung nach Anspruch 1, in der R<sup>1</sup> einen C<sub>2-5</sub>-Alkylrest bedeutet.
3. Verbindung nach Anspruch 1, in der R<sup>1</sup> einen C<sub>3-5</sub>-Alkenylrest bedeutet.
- 5 4. Verbindung nach Anspruch 1, in der R<sup>1</sup> eine Cyclopropylmethylgruppe bedeutet.
5. Verbindung nach Anspruch 1, in der R<sup>1</sup> eine Gruppe CF<sub>3</sub>, CH<sub>2</sub>CF<sub>3</sub> oder CF<sub>2</sub>CHFCF<sub>3</sub> bedeutet.
6. Verbindung nach einem der Ansprüche 1 bis 5, in der R<sup>2</sup> ein Wasserstoffatom oder eine Hydroxygruppe bedeutet.
- 10 7. Verbindung nach einem der Ansprüche 1 bis 5, in der R<sup>2</sup> eine Phenylgruppe oder einen C<sub>1-4</sub>-Alkylrest bedeutet.
8. Verbindung nach einem der Ansprüche 1 bis 5, in der R<sup>2</sup> eine Mercaptogruppe oder einen C<sub>1-4</sub>-Alkylthiorest bedeutet.
- 15 9. Verbindung nach einem der Ansprüche 1 bis 8, in der R<sup>3</sup> ein Wasserstoffatom bedeutet.
- 10 10. Verbindung nach einem der Ansprüche 1 bis 8, in der R<sup>3</sup> eine Nitro- oder Cyanogruppe oder einen Rest CONR<sup>4</sup>R<sup>5</sup> oder SO<sub>2</sub>NR<sup>4</sup>R<sup>5</sup> bedeutet.
11. Verbindung nach einem der Ansprüche 1 bis 8, in der R<sup>3</sup> ein Halogenatom, einen C<sub>1-4</sub>-Alkanoylamino-, C<sub>1-4</sub>-Alkoxy- oder C<sub>1-4</sub>-Alkylrest bedeutet.
- 25 12. Verbindung nach Anspruch 1, nämlich  
 2-(2-[2,2,2-Trifluoroethoxy]phenyl)purin-6-on,  
 2-(2-Cyclopropylmethoxyphenyl)purin-6-on,  
 2-(2-Cyclopropylmethoxyphenyl)purin-6,8-dion,  
 2-(2-Benzoyloxyphenyl)purin-6,8-dion,  
 30 2-(2-Propoxyphenyl)-8-trifluormethylpurin-6-on,  
 2-(2-Propoxyphenyl)-8-phenylpurin-6-on,  
 2-(2-Propoxyphenyl)-8-methylpurin-6-on,  
 2-(2-Propoxyphenyl)-8-mercaptapurin-6-on,  
 35 2-(2-Propoxyphenyl)-8-methylthiopurin-6-on,  
 2-(2-Propoxyphenyl)-8-aminopurin-6-on,  
 2-(2-Propoxy-5-nitrophenyl)purin-6-on,  
 2-(2-Propoxy-5-aminophenyl)purin-6-on,  
 2-(2-Propoxy-5-acetamidophenyl)purin-6-on,  
 40 2-(2-Propoxy-4-methoxyphenyl)purin-6-on,  
 2-(2-Propoxy-5-methoxyphenyl)purin-6-on,  
 2-(2-Propoxy-5-chlorphenyl)purin-6-on,  
 2-(2-Propoxy-4-methylphenyl)purin-6-on,  
 2-(2-Propoxy-5-fluorphenyl)purin-6-on,  
 45 2-(2-Propoxy-5-dimethylsulfamoylphenyl)purin-6-on,  
 2-(2-Propoxy-5-methylsulfamoylphenyl)purin-6-on,  
 2-(2-Propoxy-5-sulfamoylphenyl)purin-6-on,  
 2-(2-Propoxy-4-methylthiophenyl)purin-6-on,  
 2-(2-Propoxy-5-cyanophenyl)purin-6-on  
 50 oder 2-(2-Propoxy-5-carbamoylphenyl)purin-6-on  
 oder ein pharmazeutisch verträgliches Salz davon.
13. Verbindung nach einem der Ansprüche 1 bis 12 zur Verwendung als Medikament.
- 55 14. Arzneimittel, umfassend eine Verbindung nach einem der Ansprüche 1 bis 12 und einen pharmazeutisch verträglichen Träger.

15. Verfahren zur Herstellung einer Verbindung der Formel (1) oder eines pharmazeutisch verträglichen Salzes davon nach Anspruch 1; umfassend

- a) für Verbindung n, in denen R<sup>2</sup> ein Wasserstoffatom, einen C<sub>1-4</sub>-Alkylrest, eine Phenylgruppe oder eine Gruppe CF<sub>3</sub> bedeutet, die Umsetzung einer Verbindung der Formel (2):



(2),

in der R¹ die in Anspruch 1 angegebene Bedeutung hat und R⁶ ein nach Anspruch 1 definierter Rest R³ oder eine Vorstufe davon ist, mit einer Verbindung der Formel R⁷COL oder einem chemischen Äquivalent davon, wobei R⁷ ein Wasserstoffatom, einen C<sub>1-4</sub>-Alkylrest, eine Phenyl- oder eine Gruppe CF<sub>3</sub> bedeutet und L eine Abgangsgruppe bedeutet,

b) für Verbindungen, in denen R² eine Hydroxygruppe bedeutet, die Umsetzung einer Verbindung der vorstehend definierten Formel (2) mit einem Carbonylierungsmittel,

c) für Verbindungen, in denen R² eine Mercaptogruppe bedeutet, die Umsetzung einer Verbindung der vorstehend definierten Formel (2) mit Thioharnstoff,

d) für Verbindungen, in denen R² einen C<sub>1-4</sub>-Alkylthiorest bedeutet, die Umsetzung einer Verbindung der vorstehend definierten Formel (2) mit Thioharnstoff und danach mit einem C<sub>1-4</sub>-Alkylhalogenid,


e) für Verbindungen, in denen R² eine Aminogruppe bedeutet, die Umsetzung einer Verbindung der vorstehend definierten Formel (2) mit Cyanbromid,

und danach, falls notwendig,

- die Umwandlung eines Rests R⁶ in einen Rest R³,
- gegebenenfalls Bildung eines pharmazeutisch verträglichen Salzes.

**Patentansprüche für folgende Vertragsstaaten : ES, GR**

1. Verfahren zur Herstellung einer Verbindung der Formel (1):



(1),

oder eines pharmazeutisch verträglichen Salzes davon, wobei

R¹ einen C<sub>1-6</sub>-Alkyl-, C<sub>2-6</sub>-Alkenyl-, C<sub>3-5</sub>-Cycloalkyl-C<sub>1-4</sub>-alkyl-, Phenyl-C<sub>1-4</sub>-alkyl- oder einen mit 1 bis 6 Fluoratomen substituierten C<sub>1-4</sub>-Alkylrest bedeutet,

R² ein Wasserstoffatom, eine Hydroxygruppe, einen C<sub>1-4</sub>-Alkylrest, eine Phenyl- oder Mercaptogruppe, einen C<sub>1-4</sub>-Alkylthiorest, eine Gruppe CF<sub>3</sub> oder eine Aminogruppe bedeutet,


R³ ein Wasserstoffatom, eine Nitro- oder Aminogruppe, einen C<sub>1-4</sub>-Alkanoylamino-, C<sub>1-4</sub>-Alkoxy- oder C<sub>1-4</sub>-Alkylrest, ein Halogenatom, einen Rest -SO<sub>2</sub>NR⁴R⁵ oder

-CONR⁴R⁵, eine Cyanogruppe oder einen Rest -C<sub>1-4</sub>-AlkylS(O)<sub>n</sub> bedeutet,

R⁴ und R⁵ unabhängig voneinander ein Wasserstoffatom oder einen C<sub>1-4</sub>-Alkylrest bedeuten, und n 0, 1 oder 2 ist,

mit der Maßgabe, daß R³ kein Wasserstoffatom ist, wenn R¹ einen C<sub>1-6</sub>-Alkyl- oder C<sub>2-6</sub>-Alkenylrest

bedeutet und  $R^2$  ein Wasserstoffatom oder eine Hydroxygruppe bedeutet, wobei das Verfahren umfaßt:  
 a) für Verbindungen, in denen  $R^2$  ein Wasserstoffatom, einen  $C_1-4$ -Alkylrest, eine Phenylgruppe oder eine Gruppe  $CF_3$  bedeutet, die Umsetzung einer Verbindung der Formel (2):



(2),

in der  $R^1$  wie vorstehend definiert und  $R^6$  ein wie vorstehend definierter Rest  $R^3$  oder eine Vorstufe davon ist, mit einer Verbindung der Formel  $R^7COL$  oder einem chemischen Äquivalent davon, wobei  $R^7$  ein Wasserstoffatom, einen  $C_1-4$ -Alkylrest, eine Phenyl- oder eine Gruppe  $CF_3$  bedeutet und  $L$  eine Abgangsgruppe bedeutet,

b) für Verbindungen, in denen  $R^2$  eine Hydroxygruppe bedeutet, die Umsetzung einer Verbindung der vorstehend definierten Formel (2) mit einem Carbonylierungsmittel,

c) für Verbindungen, in denen  $R^2$  eine Mercaptogruppe bedeutet, die Umsetzung einer Verbindung der vorstehend definierten Formel (2) mit Thioharnstoff,

d) für Verbindungen, in denen  $R^2$  einen  $C_1-4$ -Alkylthioest bedeutet, die Umsetzung einer Verbindung der vorstehend definierten Formel (2) mit Thioharnstoff und danach mit einem  $C_1-4$ -Alkylhalogenid,

e) für Verbindungen, in denen  $R^2$  eine Aminogruppe bedeutet, die Umsetzung einer Verbindung der vorstehend definierten Formel (2) mit Cyanbromid,

und danach, falls notwendig,

- die Umwandlung eines Rests  $R^6$  in einen Rest  $R^3$ ,
- gegebenenfalls Bildung eines pharmazeutisch verträglichen Salzes.

2. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung, in der  $R^1$  einen  $C_2-5$ -Alkylrest bedeutet.

3. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung, in der  $R^1$  einen  $C_3-5$ -Alkenylrest bedeutet.

4. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung, in der  $R^1$  eine Cyclopropylmethylgruppe bedeutet.

5. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung, in der  $R^1$  eine Gruppe  $CF_3$ ,  $CH_2CF_3$  oder  $CF_2CHFCF_3$  bedeutet.

6. Verfahren nach einem der Ansprüche 1 bis 5 zur Herstellung einer Verbindung, in der  $R^2$  ein Wasserstoffatom oder eine Hydroxygruppe bedeutet.

7. Verfahren nach einem der Ansprüche 1 bis 5 zur Herstellung einer Verbindung, in der  $R^2$  eine Phenylgruppe oder einen  $C_1-4$ -Alkylrest bedeutet.

8. Verfahren nach einem der Ansprüche 1 bis 5 zur Herstellung einer Verbindung, in der  $R^2$  eine Mercaptogruppe oder einen  $C_1-4$ -Alkylthioest bedeutet.

9. Verfahren nach einem der Ansprüche 1 bis 8 zur Herstellung einer Verbindung, in der  $R^3$  ein Wasserstoffatom bedeutet.

10. Verfahren nach einem der Ansprüche 1 bis 8 zur Herstellung einer Verbindung, in der  $R^3$  eine Nitro- oder Cyanogruppe oder einen Rest  $CONR^4R^5$  oder  $SO_2NR^4R^5$  bedeutet.

11. Verfahren nach einem der Ansprüche 1 bis 8 zur Herstellung einer Verbindung, in der R<sup>3</sup> ein Halogenatom, einen C<sub>1-4</sub>-Alkylamino-, C<sub>1-4</sub>-Alkoxy- oder C<sub>1-4</sub>-Alkylrest bedeutet.

12. Verfahren nach Anspruch 1 zur Herstellung einer Verbindung, nämlich

- 5 2-(2-[2,2,2-Trifluorethoxy]phenyl)purin-6-on,  
 2-(2-Cyclopropylmethoxyphenyl)purin-6-on;  
 2-(2-Cyclopropylmethoxyphenyl)purin-6,8-dion,  
 2-(2-Benzoyloxyphenyl)purin-6,8-dion,  
 2-(2-Propoxyphenyl)-8-trifluormethylpurin-6-on,  
 10 2-(2-Propoxyphenyl)-8-phenylpurin-6-on,  
 2-(2-Propoxyphenyl)-8-methylpurin-6-on,  
 2-(2-Propoxyphenyl)-8-mercaptapurin-6-on,  
 2-(2-Propoxyphenyl)-8-methylthiopurin-6-on,  
 2-(2-Propoxyphenyl)-8-aminopurin-6-on,  
 15 2-(2-Propoxy-5-nitrophenyl)purin-6-on,  
 2-(2-Propoxy-5-aminophenyl)purin-6-on,  
 2-(2-Propoxy-5-acetamidophenyl)purin-6-on,  
 2-(2-Propoxy-4-methoxyphenyl)purin-6-on,  
 2-(2-Propoxy-5-methylphenyl)purin-6-on,  
 20 2-(2-Propoxy-5-chlorphenyl)purin-6-on,  
 2-(2-Propoxy-4-methylphenyl)purin-6-on,  
 2-(2-Propoxy-5-fluorphenyl)purin-6-on,  
 2-(2-Propoxy-5-dimethylsulfamoylphenyl)purin-6-on,  
 2-(2-Propoxy-5-methylsulfamoylphenyl)purin-6-on,  
 25 2-(2-Propoxy-5-sulfamoylphenyl)purin-6-on,  
 2-(2-Propoxy-4-methylthiophenyl)purin-6-on,  
 2-(2-Propoxy-5-cyanophenyl)purin-6-on  
 oder 2-(2-Propoxy-5-carbamoylphenyl)purin-6-on  
 oder ein pharmazeutisch verträgliches Salz davon.

13. Verfahren zur Herstellung eines Arzneimittels, umfassend die Vereinigung einer Verbindung nach einem der Ansprüche 1 bis 12 und eines pharmazeutisch verträglichen Trägers.


14. Verfahren nach Anspruch 1, wobei L in der Verbindung der Formel R<sup>7</sup>COL aus einer Hydroxygruppe, einem C<sub>1-4</sub>-Alkoxyrest, einem Halogenatom, einer Aminogruppe, einem C<sub>1-4</sub>-Alkylaminorest oder einem Rest OCOR<sup>7</sup> ausgewählt ist.

15. Verfahren nach Anspruch 1, wobei das Carbonylierungsmittel aus Harnstoff, Di(C<sub>1-4</sub>)-alkylcarbonat, C<sub>1-4</sub>-Alkylchlorformat, Phosgen, Trichlormethylchlorformat oder Carbonyldiimidazol ausgewählt ist.

#### Revendications

Revendications pour les Etats contractants suivants : AT, BE, CH, DE, FR, GB, IT, LI, LU, NL, SE

1. Composé de formule (1):



ou un sel de celui-ci, acceptable du point de vue pharmaceutique, dans lequel :

- R<sup>1</sup> est un groupe alkyle C<sub>1-6</sub>, alkényle C<sub>2-6</sub>, cycloalkyl C<sub>3-5</sub> alkyle C<sub>1-4</sub>, phénylalkyle C<sub>1-4</sub>, ou alkyle C<sub>1-4</sub> substitué par 1 à 6 groupes fluoro;


- R<sup>2</sup> est un atome d'hydrogène, un groupe hydroxy, alkyle C<sub>1-4</sub>, phényle, mercapto, alkyle C<sub>1-4</sub> thio, CF<sub>3</sub> ou amino;
- R<sup>3</sup> est un atome d'hydrogène, un groupe nitro, amino, alcanoyl C<sub>1-4</sub>, amino, alcoxy C<sub>1-4</sub>, alkyle C<sub>1-4</sub>, halo, SO<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>, CONR<sup>4</sup>R<sup>5</sup>, cyano, ou alkyle C<sub>1-4</sub> S(O)<sub>n</sub>;
- R<sup>4</sup> et R<sup>5</sup> sont indépendamment des atomes d'hydrogène ou des groupes alkyle C<sub>1-4</sub>; et
- n est égal à 0, 1, ou 2;

à condition que R<sup>3</sup> ne soit pas un atome d'hydrogène lorsque R<sup>1</sup> est un groupe alkyle C<sub>1-6</sub> ou alkenyle C<sub>2-6</sub>, et que R<sup>2</sup> est un atome d'hydrogène ou un groupe hydroxy.

2. Composé suivant la revendication 1, dans lequel R<sup>1</sup> est un groupe alkyle C<sub>2-5</sub>.
3. Composé suivant la revendication 1, dans lequel R<sup>1</sup> est un groupe alkenyle C<sub>3-5</sub>.
4. Composé suivant la revendication 1, dans lequel R<sup>1</sup> est un groupe cyclopropylméthyle.
5. Composé suivant la revendication 1, dans lequel R<sup>1</sup> est un groupe CF<sub>3</sub>, CH<sub>2</sub>CF<sub>3</sub>, ou CF<sub>2</sub>CHF<sub>2</sub>CF<sub>3</sub>.
6. Composé suivant l'une quelconque des revendications 1 à 5, dans lequel R<sup>2</sup> est un atome d'hydrogène ou un groupe hydroxy.
7. Composé suivant l'une quelconque des revendications 1 à 5, dans lequel R<sup>2</sup> est un groupe phényle ou alkyle C<sub>1-4</sub>.
8. Composé suivant l'une quelconque des revendications 1 à 5, dans lequel R<sup>2</sup> est un groupe mercapto, ou alkyl C<sub>1-4</sub> thio.
9. Composé suivant l'une quelconque des revendications 1 à 8, dans lequel R<sup>3</sup> est un atome d'hydrogène.
10. Composé suivant l'une quelconque des revendications 1 à 10, dans lequel R<sup>3</sup> est un groupe nitro, cyano, CONR<sup>4</sup>R<sup>5</sup>, SO<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>.
11. Composé suivant l'une quelconque des revendications 1 à 8, dans lequel R<sup>3</sup> est un groupe halo, alcanoyl C<sub>1-4</sub>, amino, alcoxy C<sub>1-4</sub> ou alkyle C<sub>1-4</sub>.
12. Composé suivant la revendication 1, qui est :
  - la 2-(2-[2,2,2-trifluoroéthoxy]phényl)purin-6-one,
  - la 2-(2-cyclopropylméthoxyphényl)purin-6-one,
  - la 2-(2-cyclopropylméthoxyphényl)purine-6,8-dione,
  - la 2-(2-benzoyloxyphényl)purine-6,8-dione,
  - la 2-(2-propoxyphényl)-8-trifluorométhylpurin-6-one,
  - la 2-(2-propoxyphényl)-8-phénylpurin-6-one,
  - la 2-(2-propoxyphényl)-8-méthylpurin-6-one,
  - la 2-(2-propoxyphényl)-8-mercaptapurin-6-one,
  - la 2-(2-propoxyphényl)-8-méthylthiopurin-6-one,
  - la 2-(2-propoxyphényl)-8-aminopurin-6-one,
  - la 2-(2-propoxy-5-nitrophényl)purin-6-one,
  - la 2-(2-propoxy-5-aminophényl)purin-6-one,
  - la 2-(2-propoxy-5-acétamidophényl)purin-6-one,
  - la 2-(2-propoxy-4-méthoxyphényl)purin-6-one,
  - la 2-(2-propoxy-5-méthoxyphényl)purin-6-one,
  - la 2-(2-propoxy-5-chlorophényl)purin-6-one,
  - la 2-(2-propoxy-4-méthylphényl)purin-6-one,
  - la 2-(2-propoxy-5-fluorophényl)purin-6-one,
  - la 2-(2-propoxy-5-diméthylsulfamoylphényl)purin-6-one,
  - la 2-(2-propoxy-5-méthylsulfamoylphényl)purin-6-one,
  - la 2-(2-propoxy-5-sulfamoylphényl)purin-6-one,
  - la 2-(2-propoxy-4-méthylthiophényl)purin-6-one,

- la 2-(2-propoxy-5-cyanophényl)purin-6-one, ou
  - la 2-(2-propoxy-5-carbamoylphényl)purin-6-one,
- ou un de leurs sels, acceptable du point de vue pharmaceutique.


13. Composé suivant l'une quelconque des revendications 1 à 12, pour l'usage en tant que médicament.
14. Composition pharmaceutique qui comprend un composé suivant l'une quelconque des revendications 1 à 12, et un véhicule acceptable du point de vue pharmaceutique.
15. Procédé de préparation d'un composé de formule (1) ou d'un sel de celui-ci, acceptable du point de vue pharmaceutique, suivant la revendication 1, qui comprend :
- (a) pour les composés dans lesquels  $R^2$  est un atome d'hydrogène, un groupe alkyle  $C_{1-4}$ , phényle, ou  $CF_3$ , la réaction d'un composé de formule (2):



- dans lequel  $R^1$  est tel que défini dans la revendication 1, et  $R^6$  est un groupe  $R^3$  tel que défini dans la revendication 1, ou un précurseur de celui-ci, avec un composé de formule  $R^7COL$ , ou un équivalent du point de vue chimique, dans lequel  $R^7$  est un atome d'hydrogène, un groupe alkyle  $C_{1-4}$ , phényle ou  $CF_3$ , et  $L$  est un groupe partant;
- (b) pour les composés dans lesquels  $R^2$  est un groupe hydroxy, la réaction d'un composé de formule (2) tel que défini plus haut, avec un agent de carbonylation;
- (c) pour les composés dans lesquels  $R^2$  est un groupe mercapto, la réaction d'un composé de formule (2) tel que défini plus haut, avec la thiourée;
- (d) pour les composés dans lesquels  $R^2$  est un groupe alkyl  $C_{1-4}$  thio, la réaction d'un composé de formule (2) tel que défini plus haut, avec la thiourée et ensuite, avec un halogénure d'alkyle  $C_{1-4}$ ;
- (e) pour les composés dans lesquels  $R^2$  est un groupe amino, la réaction d'un composé de formule (2) tel que défini plus haut, avec le bromure de cyanogène;
- et ensuite, si nécessaire :
- la conversion d'un groupe  $R^6$  en un groupe  $R^3$ ;
  - éventuellement, la formation d'un sel, acceptable du point de vue pharmaceutique.

#### Revendication pour les Etats contractants suivants : ES, GR

1. Procédé de préparation d'un composé de formule (1) :



ou un sel de celui-ci, acceptable du point de vue pharmaceutique, dans lequel :


- R<sup>1</sup> est un groupe alkyle C<sub>1-6</sub>, alkényle C<sub>2-6</sub>, cycloalkyl C<sub>3-5</sub>, alkyle C<sub>1-4</sub>, phénylalkyle C<sub>1-4</sub>, ou alkyle C<sub>1-4</sub> substitué par 1 à 6 groupes fluoro;
- R<sup>2</sup> est un atome d'hydrogène, un groupe hydroxy, alkyle C<sub>1-4</sub>, phényle, mercapto, alkyl C<sub>1-4</sub> thio, CF<sub>3</sub> ou amino;
- R<sup>3</sup> est un atome d'hydrogène, un groupe nitro, amino, alcanoyl C<sub>1-4</sub>, amino, alcoxy C<sub>1-4</sub>, alkyle C<sub>1-4</sub>, halo, SO<sub>2</sub>NR<sup>4</sup>R<sup>5</sup>, CONR<sup>4</sup>R<sup>5</sup>, cyano, ou alkyle C<sub>1-4</sub> S(O)<sub>n</sub>;
- R<sup>4</sup> et R<sup>5</sup> sont indépendamment des atomes d'hydrogène ou des groupes alkyle C<sub>1-4</sub>; et
- n est égal à 0, 1, ou 2;

à condition que R<sup>3</sup> ne soit pas un atome d'hydrogène lorsque R<sup>1</sup> est un groupe alkyle C<sub>1-6</sub> ou alkényle C<sub>2-6</sub>, et que R<sup>2</sup> est un atome d'hydrogène ou un groupe hydroxy;

lequel procédé comprend :

- (a) pour les composés dans lesquels R<sup>2</sup> est un atome d'hydrogène, un groupe alkyle C<sub>1-4</sub>, phényle, ou CF<sub>3</sub>,

la réaction d'un composé de formule (2):



(2)

dans lequel R<sup>1</sup> est tel que défini plus haut, et R<sup>6</sup> est un groupe R<sup>3</sup> tel que défini plus haut, ou un précurseur de celui-ci, avec un composé de formule R<sup>7</sup>COL; ou un équivalent du point de vue chimique, dans lequel R<sup>7</sup> est un atome d'hydrogène, un groupe alkyle C<sub>1-4</sub>, phényle ou CF<sub>3</sub>, et L est un groupe partant;

- (b) pour les composés dans lesquels R<sup>2</sup> est un groupe hydroxy,

la réaction d'un composé de formule (2) tel que défini plus haut, avec un agent de carbonylation;

- (c) pour les composés dans lesquels R<sup>2</sup> est un groupe mercapto, la réaction d'un composé de formule (2) tel que défini plus haut, avec la thiourée;

- (d) pour les composés dans lesquels R<sup>2</sup> est un groupe alkyl C<sub>1-4</sub> thio, la réaction d'un composé de formule (2) tel que défini plus haut, avec la thiourée et ensuite, avec un halogénure d'alkyle C<sub>1-4</sub>;

- (e) pour les composés dans lesquels R<sup>2</sup> est un groupe amino, la réaction d'un composé de formule (2) tel que défini plus haut, avec le bromure de cyanogène;

et ensuite, si nécessaire :

- la conversion d'un groupe R<sup>6</sup> en un groupe R<sup>3</sup>;
- éventuellement, la formation d'un sel, acceptable du point de vue pharmaceutique.

2. Procédé suivant la revendication 1, pour la préparation d'un composé dans lequel R<sup>1</sup> est un groupe alkyle C<sub>1-5</sub>.

3. Procédé suivant la revendication 1, pour la préparation d'un composé dans lequel R<sup>1</sup> est un groupe alkényle C<sub>3-5</sub>.

4. Procédé suivant la revendication 1, pour la préparation d'un composé dans lequel R<sup>1</sup> est un groupe cyclopropylméthyle.

5. Procédé suivant la revendication 1, pour la préparation d'un composé dans lequel R<sup>1</sup> est un groupe CF<sub>3</sub>, CH<sub>2</sub>CF<sub>3</sub>, ou CF<sub>2</sub>CHFCF<sub>3</sub>.

6. Procédé suivant l'une quelconque des revendications 1 à 5, pour la préparation d'un composé dans lequel R<sup>2</sup> est un atome d'hydrogène ou un groupe hydroxy.

7. Procédé suivant l'une quelconque des revendications 1 à 5, pour la préparation d'un composé dans lequel R<sup>2</sup> est un groupe phényle ou alkyle C<sub>1-4</sub>.



8. Procédé suivant l'une quelconque des revendications 1 à 5, pour la préparation d'un composé dans lequel  $R^2$  est un groupe mercapto, ou alkyl  $C_{1-4}$  thio.
9. Procédé suivant l'une quelconque des revendications 1 à 8, pour la préparation d'un composé dans lequel  $R^3$  est un atome d'hydrogène.
10. Procédé suivant l'une quelconque des revendications 1 à 8, pour la préparation d'un composé dans lequel  $R^3$  est un groupe nitro, cyano  $CONR^4R^5$  ou  $SO_2NR^4R^5$ .
11. Procédé suivant l'une quelconque des revendications 1 à 8, pour la préparation d'un composé dans lequel  $R^3$  est un groupe halo, alcanoyl  $C_{1-4}$  amino, alcoxy  $C_{1-4}$  ou alkyle  $C_{1-4}$ .
12. Procédé suivant la revendication 1, pour la préparation d'un composé qui est :
- la 2-(2-[2,2,2-trifluoroéthoxy]phényl)purin-6-one,
  - la 2-(2-cyclopropylméthoxyphényl)purin-6-one,
  - la 2-(2-cyclopropylméthoxyphényl)purine-6,8-dione,
  - la 2-(2-benzoyloxyphényl)purine-6,8-dione,
  - la 2-(2-propoxyphényl)-8-trifluorométhylpurin-6-one,
  - la 2-(2-propoxyphényl)-8-phénylpurin-6-one,
  - la 2-(2-propoxyphényl)-8-méthylpurin-6-one,
  - la 2-(2-propoxyphényl)-8-mercaptapurin-6-one,
  - la 2-(2-propoxyphényl)-8-méthylthiopurin-6-one,
  - la 2-(2-propoxyphényl)-8-aminopurin-6-one,
  - la 2-(2-propoxy-5-nitrophényl)purin-6-one,
  - la 2-(2-propoxy-5-aminophényl)purin-6-one,
  - la 2-(2-propoxy-5-acétamidophényl)purin-6-one,
  - la 2-(2-propoxy-4-méthoxyphényl)purin-6-one,
  - la 2-(2-propoxy-5-méthoxyphényl)purin-6-one,
  - la 2-(2-propoxy-5-chlorophényl)purin-6-one,
  - la 2-(2-propoxy-4-méthylphényl)purin-6-one,
  - la 2-(2-propoxy-5-fluorophényl)purin-6-one,
  - la 2-(2-propoxy-5-diméthylsulfamoylphényl)purin-6-one,
  - la 2-(2-propoxy-5-méthylsulfamoylphényl)purin-6-one,
  - la 2-(2-propoxy-5-sulfamoylphényl)purin-6-one,
  - la 2-(2-propoxy-4-méthylthiophényl)purin-6-one,
  - la 2-(2-propoxy-5-cyanophényl)purin-6-one, ou
  - la 2-(2-propoxy-5-carbamoylphényl)purin-6-one,
- ou un de leurs sels, acceptable du point de vue pharmaceutique.
13. Procédé de préparation d'une composition pharmaceutique comprenant la mise en association d'un composé suivant l'une quelconque des revendications 1 à 12, et d'un véhicule, acceptable du point de vue pharmaceutique.
14. Procédé suivant la revendication 1, dans lequel L, dans le composé de formule  $R^7COL$ , est choisi parmi les groupes hydroxy, alcoxy  $C_{1-4}$ , halo, amino, alkyl  $C_{1-4}$  amino, ou  $OCOR^7$ .
15. Procédé suivant la revendication 1, dans lequel l'agent de carbonylation est choisi parmi l'urée, les carbonates de dialkyle  $C_{1-4}$ , les chloroformiates d'alkyle  $C_{1-4}$ , le phosgène, le chloroformiate de trichlorométhyle ou le carbonyle diimidazole.